

PRVPATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

SE 00 / 756

REC 27 JUN 2000

**Intyg
Certificate**

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande Astra AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 9904419-0
Patent application number

(86) Ingivningsdatum 1999-12-03
Date of filing

Stockholm, 2000-06-21

För Patent- och registreringsverket
For the Patent- and Registration Office


Görel Gustafsson

Avgift
Fee

**CERTIFIED COPY OF
PRIORITY DOCUMENT****PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

**PATENT- OCH
REGISTRERINGSVERKET
SWEDEN**

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 866 02 86
08-666 02 86

NEW USE**Field of the Invention**

- 5 This invention relates to a new use of low molecular weight thrombin inhibitors.

Background and Prior Art

- 10 Blood coagulation is the key process involved in both haemostasis (i.e. the prevention of blood loss from a damaged vessel) and thrombosis (i.e. the formation of a blood clot in a blood vessel, sometimes leading to vessel obstruction).
- 15 Coagulation is the result of a complex series of enzymatic reactions. One of the ultimate steps in this series of reactions is the conversion of the proenzyme prothrombin to the active enzyme thrombin.

- Thrombin is known to play a central role in coagulation. It activates
- 20 platelets, leading to platelet aggregation, converts fibrinogen into fibrin monomers, which polymerise spontaneously into fibrin polymers, and activates factor XIII, which in turn crosslinks the polymers to form insoluble fibrin. Furthermore, thrombin activates factor V and factor VIII leading to a "positive feedback" generation of thrombin from
- 25 prothrombin.

Effective inhibitors of thrombin are thus known, and/or are expected, to be useful as anticoagulants and therefore useful in the therapeutic treatment of thrombosis and related disorders.

- 5 The early development of low molecular weight inhibitors of thrombin has been described by Claesson in Blood Coagul. Fibrinol. (1994) 5, 411. Low molecular weight thrombin inhibitors have been described more recently in US Patent N° 4,346,078; International Patent Applications WO 93/11152, WO 93/18060, WO 93/05069, WO 94/20467, WO 94/29336,
10 WO 95/35309, WO 95/23609, WO 96/03374, WO 96/06832, WO 96/06849, WO 96/25426, WO 96/32110, WO 97/01338, WO 97/02284, WO 97/15190, WO 97/30708, WO 97/40024, WO 97/46577, WO 98/06740, WO 97/49404, WO 97/11693, WO 97/24135, WO 97/47299, WO 98/01422 and WO 98/57932; and European Patent Applications 648
15 780, 468 231, 559 046, 641 779, 185 390, 526 877, 542 525, 195 212, 362 002, 364 344, 530 167, 293 881, 686 642, 669 317, 601 459 and 623 596.

In particular, international patent application WO 94/29336 discloses a
20 group of compounds, including $\text{HOOC-CH}_2\text{-(R)Cgl-Aze-Pab-H}$, which is also known as melagatran (see Example 1 of WO 94/29336, and the list of abbreviations in this document). International Patent Application WO 97/23499 discloses prodrugs of *inter alia* melagatran.

- 25 None of the above-mentioned documents disclose or suggest the administration of an active thrombin inhibitor in conjunction with a prodrug of that thrombin inhibitor, or indeed in conjunction with a prodrug of any thrombin inhibitor.

Deep venous thrombosis (DVT) and pulmonary embolism (PE) are major health problems, which may give rise to serious outcomes. In particular, PE may be fatal, or may result in the development of pulmonary hypertension and heart failure from recurrent embolism. DVT may result in post-thrombotic venous insufficiency and ulcers in the affected part of the body (e.g. leg). Both are common conditions, which have a great impact on worldwide healthcare costs.

There is a considerable incidence of DVT and PE following orthopaedic surgery. For example, in patients undergoing total hip replacement, the incidence of DVT in the absence of thromboprophylaxis may be as high as 45 to 57%. Further, the incidence of proximal DVT may be between 23 and 36%, and that of fatal PE, 0.34 to 6%. In patients undergoing total knee replacement in the absence of thromboprophylaxis, the postoperative incidence of DVT is between 40 and 84%, of proximal DVT is between 9 and 20%, and of fatal PE is between 0.2 and 0.7%. In patients undergoing general surgery in the absence of thromboprophylaxis, the postoperative incidence of DVT is about 25%. (Reference: Chest (1998) 114, 531S to 560S.)

20

Low-dose, subcutaneous (s.c.) unfractionated heparin is the most widely used current prophylactic treatment for venous thromboembolism resulting from orthopaedic and general surgery. The incidence of DVT after total hip replacement has been shown to be reduced (see Chest reference above).

25

The use of low-molecular weight heparin (LMWH) in the prophylaxis of DVT following total hip and knee replacement operations has been shown to further the reduce incidence (when compared to low dose unfractionated

heparin), without a concomitant increase in bleeding (see Chest reference above).

However, prolonged treatment with heparins has been shown to give rise
5 to an increased risk of osteoporosis. Heparins may also give rise to
"heparin-induced thrombocytopenia" (HIT), are dependent on the plasma
level of the endogenous thrombin inhibitor, antithrombin, and do not
inactivate clot-bound thrombin.

10 Oral anticoagulants, such as warfarin (a vitamin K antagonist), has also
been shown to be effective in reducing DVT after major surgery (see
Chest reference above). However, due to the risk of bleeding, and the
need for frequent laboratory control, the use of this substance is generally
reserved for high risk patients, and/or for long term use. Vitamin K
15 antagonists also demonstrate a notable risk of interaction with other drugs
and certain foods, and their use requires monitoring of the patient's blood
coagulation status.

Antiplaquet agents, such as aspirin, have been shown to have limited
20 efficacy in preventing DVT (see Chest reference above).

Comparative clinical studies carried out during the course of total hip
replacement operations have shown that subcutaneous administration of
the thrombin inhibitor hirudin is superior to unfractionated heparin and
25 LMWH in reducing the frequency of total and proximal DVT with no
corresponding increase in bleeding (see Eriksson *et al* in Lancet, 347, 635
(1996) and J. Bone Joint. Surg., Sep., 11 (1996)). However, hirudin is
expensive and has an immunogenic potential.

Thus, there is a need for effective treatments of thrombotic conditions such as DVT.

Disclosure of the Invention

5

We have found, surprisingly, that administration of a low molecular weight thrombin inhibitor in conjunction with a prodrug of a (or a prodrug of that) thrombin inhibitor gives rise to a notable anticoagulant effect.

10 According to a first aspect of the invention there is provided a kit of parts comprising:

(a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; and

(b) a pharmaceutical formulation including a prodrug of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative of that prodrug, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier,

20 which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

It is preferred that the prodrug of component (b) is a prodrug of the active low molecular weight thrombin inhibitor of component (a).

25

According to a further aspect of the invention, there is provided a method of making a kit of parts as defined herein, which method comprises bringing a component (a), as defined above, into association with a

component (b), as defined above, thus rendering the two components suitable for administration in conjunction with each other.

By bringing the two components "into association with" each other, we
5 include that components (a) and (b) may be:

- (i) provided as separate formulations (i.e. independently of one another), which are subsequently brought together for use in conjunction with each other in combination therapy; or
- (ii) packaged and presented together as separate components of a
10 "combination pack" for use in conjunction with each other in combination therapy.

Thus, there is further provided a kit of parts comprising:

- (1) one of components (a) and (b) as defined herein; together with
- 15 (2) instructions to use that component in conjunction with the other of the two components.

The kits of parts defined herein may comprise more than one formulation including an appropriate quantity/dose of thrombin inhibitor, and/or more
20 than one formulation including an appropriate quantity/dose of respective prodrug, in order to provide for repeat dosing. If more than one formulation (comprising thrombin inhibitor or prodrug) is present, such formulations may be the same, or may be different in terms of the dose of thrombin inhibitor/prodrug, chemical composition and/or physical form.

25

A further aspect of the invention provides a method of treatment of a condition in which inhibition of thrombin is required or desired, which comprises administration of:

- (a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; in conjunction with
 - 5 (b) a pharmaceutical formulation including a prodrug of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative of that prodrug, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier,
- to a patient suffering from, or susceptible to, such a condition.

10

For the avoidance of doubt, as used herein, the term "treatment" includes therapeutic and/or prophylactic treatment.

"Pharmaceutically acceptable derivatives" of thrombin inhibitors and
 15 prodrugs includes salts (e.g. pharmaceutically acceptable non-toxic organic or inorganic acid addition salts) and solvates.

By "administration in conjunction with", we include that respective formulations comprising thrombin inhibitor and/or prodrug are
 20 administered, sequentially, separately and/or simultaneously, over the course of treatment of the relevant condition, which condition may be acute or chronic. Preferably, the term includes that the two formulations are administered (optionally repeatedly) sufficiently closely in time for there to be a beneficial effect for the patient, that is greater, over the
 25 course of the treatment of the relevant condition, than if either of the two formulations are administered (optionally repeatedly) alone, in the absence of the other formulation, over the same course of treatment. Determination of whether a combination provides a greater beneficial effect in respect of, and over the course of treatment of, a particular

condition, will depend upon the condition to be treated or prevented, but may be achieved routinely by the skilled person.

Thus, the term "in conjunction with" includes that one or other of the two
5 formulations may be administered (optionally repeatedly) prior to, after,
and/or at the same time as, administration with the other component.
When used in this context, the terms "administered simultaneously" and
"administered at the same time as" include that individual doses of
thrombin inhibitor and prodrug are administered within 48 hours (e.g. 24
10 hours) of each other.

Components (a) and (b) as described herein may also be presented (i.e.
formulated) as a combined preparation (i.e. presented as a single
formulation including low molecular thrombin inhibitor and prodrug).

15

Thus, there is further provided a pharmaceutical formulation including a
low molecular weight thrombin inhibitor (or a pharmaceutically acceptable
derivative thereof) and a prodrug of a low molecular weight thrombin
inhibitor (or a pharmaceutically acceptable derivative of that prodrug), in
20 admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

The term "low molecular weight thrombin inhibitor" will be understood
by those skilled in the art. The term may also be understood to include
any composition of matter (e.g. chemical compound) which inhibits
25 thrombin to an experimentally determinable degree in *in vivo* and/or in *in vitro*
tests, and which possesses a molecular weight of below 2,000,
preferably below 1,000.

Preferred low molecular weight thrombin inhibitors include low molecular weight peptide-based, amino acid-based, and/or peptide analogue-based, thrombin inhibitors.

- 5 The term "low molecular weight peptide-based, amino acid-based, and/or peptide analogue-based, thrombin inhibitors" will be well understood by one skilled in the art to include low molecular weight thrombin inhibitors with one to four peptide linkages, and includes those described in the review paper by Claesson in Blood Coagul. Fibrin. (1994) 5, 411, as well as those
10 disclosed in US Patent N° 4,346,078; International Patent Applications WO 93/11152, WO 93/18060, WO 93/05069, WO 94/20467, WO 94/29336, WO 95/35309, WO 95/23609, WO 96/03374, WO 96/06832, WO 96/06849, WO 96/25426, WO 96/32110, WO 97/01338, WO 97/02284, WO 97/15190, WO 97/30708, WO 97/40024, WO 97/46577, WO
15 98/06740, WO 97/49404, WO 97/11693, WO 97/24135, WO 97/47299, WO 98/01422 and WO 98/57932; and European Patent Applications 648 780, 468 231, 559 046, 641 779, 185 390, 526 877, 542 525, 195 212, 362 002, 364 344, 530 167, 293 881, 686 642, 669 317, 601 459 and 623 596, the disclosures in all of which documents are hereby incorporated by
20 reference.

- Preferred low molecular weight peptide-based thrombin inhibitors include HOOC-CH₂-(R)Cha-Pic-Nag-H (known as inogatran; see International Patent Application WO 93/11152 and the list of abbreviations therein) and,
25 especially, HOOC-CH₂-(R)Cgl-Aze-Pab-H (known as melagatran; see International Patent Application WO 94/29336 and the list of abbreviations therein).

The term "prodrug" of a low molecular weight thrombin inhibitor includes any compound that, following oral or parenteral administration, is metabolised *in vivo* to form a low molecular weight thrombin inhibitor (as defined herein), in an experimentally-detectable amount, and within a
5 predetermined time (e.g. within a dosing interval of between 6 and 24 hours (i.e. once to four times daily)), following oral or parenteral administration. Prodrugs of the thrombin inhibitor melagatran that may be mentioned include those disclosed in international patent application WO 97/23499. Preferred prodrugs are those of the formula $R^1O_2C-CH_2-$
10 $(R)Cgl-Aze-Pab-OH$ (see the list of abbreviations in WO 97/23499), wherein R^1 represents linear or branched C_{1-6} alkyl (e.g. C_{1-4} alkyl, especially methyl, propyl and, particularly, ethyl) and the OH group replaces one of the amidino hydrogens in Pab.

15 The term "condition in which inhibition of thrombin is required or desired" will be understood by those skilled in the art to include the following:

The treatment and/or prophylaxis of thrombosis and hypercoagulability in
20 blood and tissues of animals including man. It is known that hypercoagulability may lead to thrombo-embolic diseases. Conditions associated with hypercoagulability and thrombo-embolic diseases which may be mentioned include inherited or acquired activated protein C resistance, such as the factor V-mutation (factor V Leiden), and inherited
25 or acquired deficiencies in antithrombin III, protein C, protein S, heparin cofactor II. Other conditions known to be associated with hypercoagulability and thrombo-embolic disease include circulating antiphospholipid antibodies (Lupus anticoagulant), homocysteinemi, heparin induced thrombocytopenia and defects in fibrinolysis.

The treatment of conditions where there is an undesirable excess of thrombin without signs of hypercoagulability, for example in neurodegenerative diseases such as Alzheimer's disease.

5

Particular disease states which may be mentioned include the therapeutic and/or prophylactic treatment of venous thrombosis (e.g. DVT) and pulmonary embolism, arterial thrombosis (e.g. in myocardial infarction, unstable angina, thrombosis-based stroke and peripheral arterial
10 thrombosis) and systemic embolism usually from the atrium during arterial fibrillation or from the left ventricle after transmural myocardial infarction, or caused by congestive heart failure; prophylaxis of re-occlusion (ie thrombosis) after thrombolysis, percutaneous trans-luminal angioplasty (PTA) and coronary bypass operations; the prevention of re-
15 thrombosis after microsurgery and vascular surgery in general.

Further indications include the therapeutic and/or prophylactic treatment of disseminated intravascular coagulation caused by bacteria, multiple trauma, intoxication or any other mechanism; anticoagulant treatment
20 when blood is in contact with foreign surfaces in the body such as vascular grafts, vascular stents, vascular catheters, mechanical and biological prosthetic valves or any other medical device; and anticoagulant treatment when blood is in contact with medical devices outside the body such as during cardiovascular surgery using a heart-lung machine or in
25 haemodialysis; the therapeutic and/or prophylactic treatment of idiopathic and adult respiratory distress syndrome, pulmonary fibrosis following treatment with radiation or chemotherapy, septic shock, septicemia, inflammatory responses, which include, but are not limited to, edema, acute or chronic atherosclerosis such as coronary arterial disease, cerebral

arterial disease, peripheral arterial disease, reperfusion damage, and restenosis after percutaneous trans-luminal angioplasty (PTA).

Preferred conditions include thrombosis, especially DVT, including distal
5 and proximal DVT. The present invention finds particular utility in the prophylactic treatment of DVT resulting from surgery, such as gastrointestinal, or orthopaedic, surgery (e.g. hip or knee replacement). This includes DVT resulting from immobilisation after surgery.

10 In accordance with the invention, thrombin inhibitors, prodrugs of thrombin inhibitors, and derivatives of either, may be administered orally, intravenously, subcutaneously, buccally, rectally, dermally, nasally, tracheally, bronchially, topically, by any other parenteral route, or *via* inhalation, in the form of a pharmaceutical preparation comprising the
15 thrombin inhibitor or prodrug in a pharmaceutically acceptable dosage form. Depending on the disorder, and the patient, to be treated, as well as the route of administration, the compositions may be administered at varying doses.

20 Preferred modes of delivery are systemic. For melagatran and derivatives thereof, preferred modes of administration are parenteral, more preferably intravenous, and especially subcutaneous. For prodrugs of melagatran, preferred modes of administration are oral.

25 In the therapeutic treatment of mammals, and especially humans, thrombin inhibitors, prodrugs of thrombin inhibitors, and derivatives of either, may be administered alone, but will generally be administered as a pharmaceutical formulation in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, which may be selected with due

regard to the intended route of administration and standard pharmaceutical practice.

Suitable formulations for use in administering thrombin inhibitors are
 5 known in the art, and include those known from US Patent N° 4,346,078;
 International Patent Applications WO 93/11152, WO 93/18060, WO
 93/05069, WO 94/20467, WO 94/29336, WO 95/35309, WO 95/23609,
 WO 96/03374, WO 96/06832, WO 96/06849, WO 96/25426, WO
 96/32110, WO 97/01338, WO 97/02284, WO 97/15190, WO 97/30708,
 10 WO 97/40024, WO 97/46577, WO 98/06740, WO 97/49404, WO
 97/11693, WO 97/24135, WO 97/47299, WO 98/01422 and WO
 98/57932; and European Patent Applications 648 780, 468 231, 559 046,
 641 779, 185 390, 526 877, 542 525, 195 212, 362 002, 364 344, 530
 167, 293 881, 686 642, 669 317, 601 459 and 623 596, the disclosures in
 15 all of which documents are hereby incorporated by reference.

Suitable formulations for use with melagatran, derivatives and prodrugs
 thereof are described in the literature, for example as described in *inter*
alia international patent applications WO 94/29336, WO 96/14084, WO
 20 96/16671, WO 97/23499, WO 97/39770, WO 97/45138 and WO
 98/16252, the disclosures in which documents are hereby incorporated by
 reference. Otherwise, the preparation of suitable formulations may be
 achieved non-inventively by the skilled person using routine techniques.

25 The amounts of thrombin inhibitor, prodrug, or derivative of either, in the
 formulation will depend on the severity of the condition, and on the
 patient, to be treated, as well as the compound(s) which is/are employed,
 but may be determined non-inventively by the skilled person.

Suitable doses of thrombin inhibitors, prodrugs and derivatives of either, in the therapeutic and/or prophylactic treatment of mammalian, especially human, patients may be determined routinely by the medical practitioner or other skilled person, and include the respective doses discussed in the prior art documents disclosing thrombin inhibitors that are mentioned
5 hereinbefore, the disclosures in which are incorporated by reference.

In the case of melagatran, suitable doses of active compound, prodrugs and derivatives thereof, in the therapeutic and/or prophylactic treatment of mammalian, especially human, patients include those which give a mean
10 plasma concentration of up to 5 $\mu\text{mol/L}$, for example in the range 0.001 to 5 $\mu\text{mol/L}$ over the course of treatment of the relevant condition.

In any event, the physician, or the skilled person, will be able to determine the actual dosage which will be most suitable for an individual
15 patient, which is likely to vary with the condition that is to be treated, as well as the age, weight, sex and response of the particular patient to be treated. The above-mentioned dosages are exemplary of the average case; there can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.
20

The sequence in which the formulations comprising thrombin inhibitor, and prodrug, may be administered (i.e. whether, and at what point, sequential, separate and/or simultaneous administration takes place) may
25 be determined by the physician or skilled person. For example, the sequence may depend upon many factors that will be evident to the skilled person, such as whether, at any time during the course or period of treatment, one or other of the formulations cannot be administered to the patient for practical reasons (e.g. the patient is unconscious and thus

unable to take an oral formulation comprising either thrombin inhibitor or prodrug).

For example, in the treatment of thrombosis (e.g. DVT) resulting from surgery, such as gastrointestinal, or orthopaedic, surgery, and when the active thrombin inhibitor is melagatran, it is preferred that the formulation comprising melagatran is administered parenterally within two days (e.g. within 24 hours) of surgery (either prior to or after surgery), and particularly immediately prior to (e.g. within 2 hours), and/or within up to 12 hours after surgery (e.g. at least one hour after surgery), and thereafter for up to between 3 and 7 (e.g. between 0 and 2, such as between 1 and 2) days after that surgery, and that the formulation comprising prodrug is administered orally within 7 days following that surgery (preferably once administration of melagatran has been terminated) for up to e.g. between 11 and 40 days, preferably 9 days, more preferably up to 8 days.

The method described herein may have the advantage that, in the treatment of conditions in which inhibition of thrombin is required or desired, it may be more convenient for the physician and/or patient than, be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, produce fewer side effects than, or that it may have other useful pharmacological properties over, similar methods known in the prior art for the treatment of such conditions.

The invention is illustrated, but in no way limited, by the following example.

Example 1

Clinical Trial - Melagatran and EtOOC-CH₂-(R)Cgl-Aze-Pab-OH
Combination Therapy

- 5 A controlled, randomised, parallel group, Swedish multi-centre pilot study was carried out. The study was open with regard to the drugs under evaluation but was blind for the patients, all personnel at the study sites, and for the person monitoring the experiments with regard to the doses of melagatran and the prodrug of melagatran, EtOOC-CH₂-(R)Cgl-Aze-Pab-
10 OH (P; see WO 97/23499).

Dalteparin (Fragmin®; Pharmacia-Upjohn) was used as a reference compound.

- 15 Patients scheduled for primary elective total hip or knee replacement were eligible for inclusion, and were randomly selected into one of three groups, each to receive different doses of melagatran and P, or dalteparin. In all, 135 patients were included in the study, of which 105 patients could be used for evaluation with respect to thromboembolic events using central
20 assessment of locally performed phlebograms.

- About 32 patients in each treatment group were evaluated according to the protocol. A stratified randomisation, by centre and type of surgery, was used to ensure that approximately equal numbers of patients were given
25 each of the drugs under evaluation at all participating centres (in all six centres were used) for both types of surgery (hip or knee). Each centre received study drugs in blocks of four, separately for hips and knees. Within each block, the order of the study drugs was randomised.

The following formulations were used in the study:

Melagatran - 5, 10 or 20 mg/mL in aqueous saline solution.

- 5 P - appropriate weight (see below) in a tablet also comprising 59 to 63 mg corn starch, 115 mg microcrystalline cellulose and 2 mg sodium stearyl fumarate.

The following doses of melagatran and P were used in the study:

10

Treatment A - s.c. melagatran (1 mg) b.i.d. for 2 days, followed by oral administration of P (6 mg) b.i.d. for 6 to 9 days.

- 15 Treatment B - s.c. melagatran (2 mg) b.i.d. for 2 days, followed by an oral administration of P (12 mg) b.i.d. for 6 to 9 days.

Treatment C - s.c. melagatran (4 mg) b.i.d. for 2 days, followed by an oral administration of P (24 mg) b.i.d. for 6 to 9 days.

- 20 The patients receiving melagatran and P received treatment on the day of surgery. The patient received the first injection after induction of anaesthesia immediately before surgery. For knee-patients, the pre-operative melagatran injection was given before tourniquets were applied. The second injection was given in the evening the same day. The patient
25 received one melagatran injection in the morning and one in the evening over the next 24 hours, until oral administration of P, twice daily, started. The first oral dose of P was always taken in the morning. Thus, the total treatment period comprised was between 8 and 11 days.

Treatment D - dalteparin (Fragmin®): one s.c. injection of 5000 U during the evening of the day before surgery, continuing with one s.c. injection every evening over a treatment period of 8 to 11 days.

- 5 The plasma concentrations of melagatran were recorded.

The results of the trial, in terms of the frequencies of thromboembolism after hip or knee surgery, are tabulated below:

	Treatment A		Treatment B		Treatment C		Treatment D	
	(n)	(%)	(n)	(%)	(n)	(%)	(n)	(%)
Outcome	6/29	21	6/24	25	4/24	16	5/27	19

10

These data show that a combination of subcutaneously administered melagatran and orally administered P is effective in preventing DVT after orthopaedic surgery.

Claims

1. A kit of parts comprising:
 - 5 (a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; and
 - (b) a pharmaceutical formulation including a prodrug of a low
10 molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative of that prodrug, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier,
which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.
- 15 2. A kit of parts as claimed in Claim 1, wherein the prodrug of component (b) is a prodrug of the thrombin inhibitor of component (a).
3. A kit of parts as claimed in Claim 1 or Claim 2, wherein components
20 (a) and (b) are suitable for sequential, separate and/or simultaneous use in the treatment of a condition in which inhibition of thrombin is required or desired.
4. A kit of parts as claimed in Claim 3, wherein the condition is deep
25 venous thrombosis.
5. A kit of parts as claimed in any one of Claims 1 to 4, wherein the thrombin inhibitor is melagatran.

6. A kit of parts as claimed in Claim 5, wherein the prodrug is of the formula



wherein R^1 represents linear or branched C_{1-6} alkyl and the OH group
5 replaces one of the amidino hydrogens in Pab.

7. A kit of parts as claimed in Claim 6, wherein R^1 represents methyl, ethyl or propyl.

10 8. A kit of parts as claimed in any one of the preceding claims, wherein the formulation comprising thrombin inhibitor, or derivative thereof, is a parenteral formulation and that comprising the prodrug, or derivative thereof, is an oral formulation.

15 9. A method of making a kit of parts as defined in any one of Claims 1 to 8, which method comprises bringing a component (a) according to any one of Claims 1 to 8, into association with a component (b) according to any one of Claims 1 to 8, thus rendering the two components suitable for administration in conjunction with each other.

20

10. A kit of parts comprising:

(1) one of components (a) and (b) as defined in any one of Claims 1 to 8; together with

(2) instructions to use that component in conjunction with the other of the
25 two components.

11. A pharmaceutical formulation including a low molecular weight thrombin inhibitor (or a pharmaceutically acceptable derivative thereof) and a prodrug of a low molecular weight thrombin inhibitor (or a

pharmaceutically acceptable derivative of that prodrug), in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier.

12. A method of treatment of a condition in which inhibition of thrombin is required or desired, which comprises administration of:

- (a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; in conjunction with
 - (b) a pharmaceutical formulation including a prodrug of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative of that prodrug, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier,
- to a patient suffering from, or susceptible to, such a condition.

15

13. A method as claimed in Claim 12 in which component (a) is administered prior to commencement of administration of component (b).

14. A method of treatment of a condition in which inhibition of thrombin is required or desired, which comprises administration of a formulation as defined in Claim 11 to a patient suffering from, or susceptible to, such a condition.

15. A method as claimed in any one of Claims 12 to 14, wherein the condition is deep venous thrombosis.

16. A method as claimed in Claim 15, wherein the thrombosis results from surgery.

17. A method as claimed in Claim 16, wherein the surgery is gastrointestinal surgery or orthopaedic surgery.

18. A method as claimed in Claim 16 or Claim 17, wherein component
5 (a) is administered parenterally prior to and/or after surgery and
component (b) is administered orally following that surgery.

19. The use of a thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in the manufacture of a medicament for the treatment or prophylaxis of a condition in which inhibition of thrombin is required or desired, which treatment or prophylaxis comprises administration of:

(a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; in conjunction with

(b) a pharmaceutical formulation including a prodrug of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative of that prodrug, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier,

20 to a patient suffering from, or susceptible to, such a condition.

ABSTRACT

According to the invention there is provided a kit of parts comprising:

- 5 (a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier; and
- 10 (b) a pharmaceutical formulation including a prodrug of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative of that prodrug, in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier,

which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other, as well as the use of such a kit of parts in the treatment of a condition in which inhibition of
15 thrombin is required or desired.